

## Summary

<b>Production Name</b>	Histone H3.3 Rabbit Polyclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,ICC/IF
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification</b>	Affinity Chromatography

## Immunogen

<b>Gene Name</b>	H3-3A
<b>Alternative Names</b>	H3F3; H3.3A
<b>Gene ID</b>	3020
<b>SwissProt ID</b>	P84243

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
<b>Molecular Weight</b>	Calculated MW: 15 kDa; Observed MW: 15 kDa

## Background

---

**Product Name: Histone H3.3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab00058**




H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


## Research Area

Epigenetics and Nuclear Signaling

## Image Data



Western blot analysis of Histone H3.3 in (1) HeLa lysates; (2) NIH/3T3 lysates using Histone H3.3 antibody.



Immunohistochemistry analysis of paraffin-embedded mouse stomach using Histone H3.3 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

## Note

For research use only.