

Summary

Production Name	PARK7/DJ1 Rabbit Polyclonal Antibody
Description	Primary antibody
Host	Rabbit
Application	WB,IHC-P,ICC/IF,FC,IP
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide
	and 50% glycerol.
Purification	Affinity Chromatography

Immunogen

Gene Name	PARK7
Alternative Names	PARK7; Protein DJ-1; Oncogene DJ1; Parkinson disease protein 7
Gene ID	11315
SwissProt ID	Q99497

Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20 FC: 1/50-
	1/100
Molecular Weight	Calculated MW: 20 kDa; Observed MW: 20 kDa


Background

Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It cooperates with Ras to increase cell transformation, it positively regulates transcription of the androgen receptor, and it may function as an indicator of oxidative stress.


Research Area

Neuroscience

Image Data



Western blot analysis of PARK7 in HeLa lysates using PARK7/DJ1 antibody.



Immunohistochemistry analysis of paraffin-embedded Human bladder cancer using PARK7 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.