

## Summary

Production Name	Cyclin E2 Rabbit Polyclonal Antibody
Description	Primary antibody
Host	Rabbit
Application	WB,IHC-P,ICC/IF,FC
Reactivity	Human, Mouse

#### Performance

Conjugation Unconjugated	
Modification Unmodified	
lsotype lgG	
Clonality Polyclonal Antibody	
Form Liquid	
Store at 4°C short term. Aliquot and store at <b>Storage</b>	t -20°C long term. Avoid freeze/thaw
cycles.	
Rabbit IgG in phosphate buffered saline , pH 7	7.4, 150mM NaCl, 0.02% sodium azide
and 50% glycerol.	
Purification Affinity Chromatography	

## Immunogen

Gene Name	CCNE2
Alternative Names	CCNE2; G1/S-specific cyclin-E2
Gene ID	9134
SwissProt ID	O96020

# Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 FC: 1/50-1/100
Molecular Weight	Calculated MW: 47 kDa; Observed MW: 47 kDa

### Background

## Product Name: Cyclin E2 Rabbit Polyclonal Antibody Catalog #: APRab00108




The human Cyclin E2 gene encodes a 404 amino acid protein that is most closely related to Cyclin E. Cyclin E2 mRNA levels peaks at the G1 / S transition. Cyclin E2 associates with Cdk2 in a functional kinase complex that is inhibited by both p27 (Kip1) and p21 (Cip1). Cyclin E2 / Cdk2 phosphorylates histone H1 in vitro. G1 cyclin E controls the initiation of DNA synthesis by activating CDK2. Abnormally high levels of cyclin E expression have frequently been observed in human cancers.


#### **Research Area**

Cell Biology

#### Image Data




Western blot analysis of Cyclin E2 in HeLa lysates using Cyclin E2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human breast carcinoma using Cyclin E2 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunofluorescence analysis of Cyclin E2 in Hela using Cyclin E2 antibody.

**Note** For research use only.