

Summary

Aryl Hydrocarbon Receptor Rabbit Polyclonal Antibody	
Primary antibody	
Rabbit	
WB,ICC/IF	
Human	

Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	IgG	
Clonality	Polyclonal Antibody	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide	
	and 50% glycerol.	
Purification	Affinity Chromatography	

Immunogen

Gene Name	AHR	
Alternative Names	Ah receptor; AhR; Class E basic helix-loop-helix protein 76; bHLHe76; AHR	
Gene ID	196	
SwissProt ID	P35869	

Application

Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200
Molecular Weight	Calculated MW: 96 kDa; Observed MW: 105 kDa




Background

The aryl hydrocarbon receptor (AhR) is a ligand activated transcription factor involved in xenobiotic metabolism, cell cycle regulation, and development in response to both endogenous and environmental signals. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues.

Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of AhR in HEK293 lysates using Aryl Hydrocarbon Receptor antibody.

Note

For research use only.