

Summary

| Production Name | DDB1 Rabbit Polyclonal Antibody |
|-----------------|---------------------------------|
| Description | Primary antibody |
| Host | Rabbit |
| Application | WB,IHC-P,ICC/IF,IP |
| Reactivity | Human,Mouse,Rat |
| | |

Performance

| Conjugation Unconjugated | |
|--|--------------------------------------|
| Modification Unmodified | |
| lsotype lgG | |
| Clonality Polyclonal Antibody | |
| Form Liquid | |
| Store at 4°C short term. Aliquot and store at Storage | t -20°C long term. Avoid freeze/thaw |
| cycles. | |
| Rabbit IgG in phosphate buffered saline , pH 7 | 7.4, 150mM NaCl, 0.02% sodium azide |
| and 50% glycerol. | |
| Purification Affinity Chromatography | |

Immunogen

| Gene Name | DDB1 |
|-------------------|--|
| Alternative Names | XPE; DDBA; XAP1; XPCE; XPE-BF; UV-DDB1 |
| Gene ID | 1642 |
| SwissProt ID | Q16531 |

Application

| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20 |
|------------------|--|
| Molecular Weight | Calculated MW: 127 kDa; Observed MW: 127 kDa |

Background

Product Name: DDB1 Rabbit Polyclonal Antibody Catalog #: APRab00331




Required for DNA repair. Binds to DDB2 to form the UV-damaged DNA-binding protein complex (the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair.

Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of DDB1 in HeLa lysates using DDB1 antibody.

Note

For research use only.