

**Product Name: RFWD3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab00385**



## Summary

<b>Production Name</b>	RFWD3 Rabbit Polyclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,ELISA
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
<b>Purification</b>	Affinity Chromatography

## Immunogen

<b>Gene Name</b>	RFWD3
<b>Alternative Names</b>	FANCW; RNF201
<b>Gene ID</b>	55159
<b>SwissProt ID</b>	Q6PCD5

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000
<b>Molecular Weight</b>	Calculated MW: 85 kDa; Observed MW: 85 kDa

## Background

E3 ubiquitin-protein ligase required for the repair of DNA interstrand cross-links (ICL) in response to DNA damage

**Product Name: RFWD3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab00385**




(PubMed:21504906, PubMed:21558276, PubMed:26474068, PubMed:28575657, PubMed:28575658). Plays a key role in RPA-mediated DNA damage signaling and repair (PubMed:21504906, PubMed:21558276, PubMed:26474068, PubMed:28575657, PubMed:28575658, PubMed:28691929). Acts by mediating ubiquitination of the RPA complex (RPA1, RPA2 and RPA3 subunits) and RAD51 at stalled replication forks, leading to remove them from DNA damage sites and promote homologous recombination (PubMed:26474068, PubMed:28575657, PubMed:28575658). Also mediates the ubiquitination of p53/TP53 in the late response to DNA damage, and acts as a positive regulator of p53/TP53 stability, thereby regulating the G1/S DNA damage checkpoint (PubMed:20173098). May act by catalyzing the formation of short polyubiquitin chains on p53/TP53 that are not targeted to the proteasome (PubMed:20173098). In response to ionizing radiation, interacts with MDM2 and enhances p53/TP53 ubiquitination, possibly by restricting MDM2 from extending polyubiquitin chains on ubiquitinated p53/TP53 (PubMed:20173098).


## Research Area

-

## Image Data




Western blot analysis of RFWD3 in 823 lysates using RFWD3 antibody.



Western blot analysis of RFWD3 in MCF-7 lysates using RFWD3 antibody. The lane on the right is blocked with the synthesized peptide.

**Product Name: RFWD3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab00385**



Western blot analysis of RFWD3 in HepG2 lysates using RFWD3 antibody.

**Note**

For research use only.