

Summary

Production Name	Phospho-Histone H3 (Thr3) Rabbit Polyclonal Antibody
Description	Primary antibody
Host	Rabbit
Application	WB,IHC-P,ICC/IF,FC
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Phosphorylated
lsotype	lgG
Clonality	Polyclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide
	and 50% glycerol.
Purification	Affinity Chromatography

Immunogen

Gene Name	H3-3A
Alternative Names	H3F3; H3.3A
Gene ID	3020
SwissProt ID	P84243

Application

Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 FC: 1/50-1/100
Molecular Weight	Calculated MW: 15 kDa; Observed MW: 15 kDa




Background

H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of Phospho-Histone H3 (Thr3) in HeLa lysates treated with FBS + Calyculin A using Phospho-Histone



Immunohistochemistry analysis of paraffin-embedded Human tonsil using Phospho-Histone H3 (T3) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.