

## Summary

ody	
Primary antibody	
WB,FC	

### Performance

Conjugation	Unconjugated	
Modification	Phosphorylated	
lsotype	IgG	
Clonality	Polyclonal Antibody	
Form	Liquid	
Storago	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
Storage	cycles.	
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide	
builei	and 50% glycerol.	
Purification	Affinity Chromatography	

#### Immunogen

Gene Name	SRC
Alternative Names	SRC; SRC1; Proto-oncogene tyrosine-protein kinase Src; Proto-oncogene c-Src; pp60c-
	src; p60-Src
Gene ID	6714
SwissProt ID	P12931

# Application

Dilution Ratio	WB: 1/500-1/1000 FC: 1/50-1/100
Molecular Weight	Calculated MW: 60 kDa; Observed MW: 60 kDa




#### Background

Src activity is regulated by tyrosine phosphorylation at two sites, but with opposing effects. While phosphorylation at Tyr416 in the activation loop of the kinase domain upregulates enzyme activity, phosphorylation at Tyr527 in the carboxyterminal tail by Csk renders the enzyme less active (2).

#### **Research Area**

**Signal Transduction** 

### **Image Data**



Western blot analysis of Phospho-Src (Y419) in A431 lysates treated with pervanadate using Phospho-Src (Tyr419) antibody.

#### Note

For research use only.