

**Product Name: Phospho-Rb (Ser807) Rabbit Polyclonal Antibody**  
**Catalog #: APRab00712**



---

## Summary

<b>Production Name</b>	Phospho-Rb (Ser807) Rabbit Polyclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-F,IHC-P,ICC/IF
<b>Reactivity</b>	Human,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phosphorylated
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA
<b>Purification</b>	Affinity Purified

## Immunogen

<b>Gene Name</b>	RB1
<b>Alternative Names</b>	RB1; Retinoblastoma-associated protein; p105-Rb; pRb; Rb; pp110
<b>Gene ID</b>	5925
<b>SwissProt ID</b>	P06400

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
<b>Molecular Weight</b>	Calculated MW: 106 kDa; Observed MW: 110 kDa

**Product Name: Phospho-Rb (Ser807) Rabbit Polyclonal Antibody**  
**Catalog #: APRab00712**




## Background

Cell cycle-dependent phosphorylation by a CDK inhibits Rb target binding and allows cell cycle progression. Rb inactivation and subsequent cell cycle progression likely requires an initial phosphorylation by cyclin D-CDK4/6 followed by cyclin E-CDK2 phosphorylation. Specificity of different CDK/cyclin complexes has been observed in vitro and cyclin D1 is required for Ser780 phosphorylation in vivo.

## Research Area

Cell Biology

## Image Data



Western blot analysis of Rb (Phospho- S807) in CHO-K1, C6 lysates using Phospho-Rb (Ser807) antibody.

## Note

For research use only.