

**Product Name: Phospho-4E BP1 (Thr70) Rabbit  
Polyclonal Antibody  
Catalog #: APRab00910**

---

## Summary

<b>Production Name</b>	Phospho-4E BP1 (Thr70) Rabbit Polyclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC-P
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phosphorylated
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
<b>Purification</b>	Affinity Purified

## Immunogen

<b>Gene Name</b>	EIF4EBP1 EIF4EBP1; Eukaryotic translation initiation factor 4E-binding protein 1; 4E-BP1; eIF4E-binding protein 1; Phosphorylated heat- and acid-stable protein regulated by insulin 1;
<b>Alternative Names</b>	PHAS-I
<b>Gene ID</b>	1978
<b>SwissProt ID</b>	Q13541

## Application

<b>Dilution Ratio</b>	IHC: 1/50-1/100
<b>Molecular Weight</b>	-

**Product Name: Phospho-4E BP1 (Thr70) Rabbit  
Polyclonal Antibody  
Catalog #: APRab00910**

---


## Background

Translation repressor protein 4E-BP1 (also known as PHAS-1) inhibits cap-dependent translation by binding to the translation initiation factor eIF4E. Hyperphosphorylation of 4E-BP1 disrupts this interaction and results in activation of cap-dependent translation. Both the PI3 kinase/Akt pathway and FRAP/mTOR kinase regulate 4E-BP1 activity.

## Research Area

Epigenetics and Nuclear Signaling

## Image Data



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using Phospho-4E BP1 (Thr70) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Sample with blocking peptide on the right.

## Note

For research use only.