

## Summary

<b>Production Name</b>	PKC delta Rabbit Polyclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-P,ICC/IF
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification</b>	Affinity Chromatography

## Immunogen

<b>Gene Name</b>	PRKCD
<b>Alternative Names</b>	Protein kinase C delta type; Tyrosine-protein kinase PRKCD; nPKC-delta
<b>Gene ID</b>	5580
<b>SwissProt ID</b>	Q05655

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
<b>Molecular Weight</b>	Calculated MW: 78 kDa; Observed MW: 78 kDa

## Background

**Product Name: PKC delta Rabbit Polyclonal Antibody**  
**Catalog #: APRab01337**




Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways.

## Research Area

Signal Transduction

## Image Data



Western blot analysis of PKC delta in HeLa lysates using PKC delta antibody.

## Note

For research use only.