

Summary

Production Name	Ferritin Heavy Chain Rabbit Polyclonal Antibody
Description	Primary antibody
Host	Rabbit
Application	ELISA
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal Antibody
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purified

Immunogen

Gene Name	FTH1
Alternative Names	FTH1; FTH; FTHL6; OK/SW-cl.84; PIG15; Ferritin heavy chain; Ferritin H subunit; Cell
	proliferation-inducing gene 15 protein
Gene ID	2495
SwissProt ID	P02794

Application

Dilution Ratio	ELISA: 1/10000
Molecular Weight	-




Background

The assembled ferritin molecule, often referred to as a nanocage, can store up to 4,500 atoms of iron. It forms a holoenzyme of ~450 kDa, consisting of 24 subunits made up of two types of polypeptide chains: ferritin heavy chain and ferritin light chain, each having unique functions. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe(II), whereas ferritin light chains promote the nucleation of ferrihydrite, enabling storage of Fe(III).

Research Area

Neuroscience

Image Data



Standard Curve for Ferritin: Capture Antibody mouse mAb (201067, Ferritin Light Chain mouse mAb) to Ferritin at 2u03bcg/ml and Detector Antibody Rabbit pAb(3010667, Ferritin rabbit pAb)to Ferritin at 0.5u03bcg/ml.

Note

For research use only.