

Summary

| Production Name | Acetyl-p53 (Lys382) Rabbit Polyclonal Antibody | |
|-----------------|--|--|
| Description | Primary antibody | |
| Host | Rabbit | |
| Application | WB,IHC-P,ELISA | |
| Reactivity | Human | |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Acetylated |
| lsotype | IgG |
| Clonality | Polyclonal Antibody |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3. |
| Purification | Affinity Purified |

Immunogen

| Gene Name | TP53 |
|-------------------|--|
| Alternative Names | TP53; P53; Cellular tumor antigen p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor |
| | suppressor p53 |
| Gene ID | 7157 |
| SwissProt ID | P04637 |

Application

| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 ELISA: 1/10000 |
|------------------|---|
| Molecular Weight | Calculated MW: 44 kDa; Observed MW: 53 kDa |




Background

Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle, specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, oligomerization and transcription activation domains.

Research Area

Cell Biology

Image Data



Western blot analysis of Acetyl-p53 (Lys382) in HepG2 lysates using Acetyl-p53 (Lys382) antibody.

Note

For research use only.