

Summary

| Production Name | GRP78 BiP Rabbit Polyclonal Antibody |
|-----------------|--------------------------------------|
| Description | Primary antibody |
| Host | Rabbit |
| Application | WB,IHC-F,IHC-P,ICC/IF,ELISA |
| Reactivity | Human, Mouse, Rat |

Performance

| Conjugation | Unconjugated | |
|--------------|--|--|
| Modification | Unmodified | |
| lsotype | IgG | |
| Clonality | Polyclonal Antibody | |
| Form | Liquid | |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw | |
| | cycles. | |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3. | |
| Purification | Affinity Purified | |

Immunogen

| Gene Name | HSPA5 |
|-------------------|---|
| | HSPA5; GRP78; 78 kDa glucose-regulated protein; GRP-78; Endoplasmic reticulum |
| Alternative Names | lumenal Ca(2+)-binding protein grp78; Heat shock 70 kDa protein 5; Immunoglobulin |
| | heavy chain-binding protein; BiP |
| Gene ID | 3309 |
| SwissProt ID | P11021 |

Application

| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ELISA: 1/10000 |
|------------------|--|
| Molecular Weight | Calculated MW: 72 kDa; Observed MW: 75 kDa |




Background

When Chinese hamster K12 cells are starved of glucose, the synthesis of several proteins, called glucose-regulated proteins (GRPs), is markedly increased. Hendershot et al. (1994) (PubMed 8020977) pointed out that one of these, GRP78 (HSPA5), also referred to as 'immunoglobulin heavy chain-binding protein' (BiP), is a member of the heat-shock protein-70 (HSP70) family and is involved in the folding and assembly of proteins in the endoplasmic reticulum (ER).

Research Area

Tags & Cell Markers

Image Data





Note

For research use only.