

Product Name: Acetyl Histone H2A (K5) Rabbit Polyclonal Antibody
Catalog #: APRab04161

Summary

Production Name	Acetyl Histone H2A (K5) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Acetyl Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	H2AFZ
Alternative Names	H2AFZ; H2AZ; Histone H2A.Z; H2A/z
Gene ID	3015.0
SwissProt ID	P0C0S5.The antiserum was produced against synthesized peptide derived from human Histone H2A around the acetylated site of Lys5. AA range:1-50

Application

Dilution Ratio	IHC 1:100-1:300 ELISA: 1:5000
Molecular Weight	14kD

Background


**Product Name: Acetyl Histone H2A (K5) Rabbit
Polyclonal Antibody
Catalog #: APRab04161**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality. [provided by RefSeq, Jul 2008],function:Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. May be involved in the formation of constitutive heterochromatin. May be required for chromosome segregation during cell division.,mass spectrometry:Monoisotopic, not modified PubMed:16457589,PTM:Acetylated on Lys-5, Lys-8 and Lys-12 during interphase. Acetylation disappears at mitosis.,PTM:Monoubiquitination of Lys-122 gives a specific tag for epigenetic transcriptional repression.,PTM:Not phosphorylated.,similarity:Belongs to the histone H2A family.,subunit:The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers. The octamer wraps approximately 147 bp of DNA. H2A or its variant H2AFZ forms an heterodimer with H2B. H2AFZ interacts with INCENP.,

Research Area

Protein_Acetylation

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Histone H2A (Acetyl-Lys5) Antibody. The picture on the right is blocked with the synthesized peptide.

**Product Name: Acetyl Histone H2A (K5) Rabbit
Polyclonal Antibody
Catalog #: APRab04161**



Note

For research use only.