



Catalog #: APRab04277

Summary

Production Name ATF-2 (phospho Thr69) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC,ELISA **Reactivity** Human,Mouse,Rat

Performance

Conjugation Unconjugated

Modification Phospho Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name ATF2

ATF2; CREB2; CREBP1; Cyclic AMP-dependent transcription factor ATF-2; cAMP-

Alternative Names dependent transcription factor ATF-2; Activating transcription factor 2; Cyclic AMP-

responsive element-binding protein 2; CREB-2; cAMP-responsive element-binding pro

Gene ID 1386.0

P15336.The antiserum was produced against synthesized peptide derived from human SwissProt ID

ATF2 around the phosphorylation site of Thr69 or 51. AA range:36-85

Application

Dilution Ratio WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000...

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: APRab04277



Molecular Weight


Background

activating transcription factor 2(ATF2) Homo sapiens This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription. This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. The encoded protein may also be involved in cell's DNA damage response independent of its role in transcriptional regulation. Several alternatively spliced transcript variants have been found for this gene [provided by RefSeq, Jan 2014caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Transcriptional activator, probably constitutive, which binds to the cAMP-responsive element (CRE) (consensus: 5'-GTGACGT[AC][AG]-3'), a sequence present in many viral and cellular promoters. Interaction with JUN redirects JUN to bind to CRES preferentially over the 12-Otetradecanoylphorbol-13-acetate response elements (TRES) as part of an ATF2-c-Jun complex., PTM: Phosphorylation of Thr-69 and Thr-71 by MAPK14 causes increased transcriptional activity. Also phosphorylated and activated by JNK., similarity: Belongs to the bZIP family., similarity: Belongs to the bZIP family. ATF subfamily., similarity: Contains 1 bZIP domain.,similarity:Contains 1 C2H2-type zinc finger.,subunit:Binds DNA as a dimer and can form a homodimer in the absence of DNA. Can form a heterodimer with JUN. Interacts with SMAD3 and SMAD4. Binds through its N-terminal region to UTF1 which acts as a coactivator of ATF2 transcriptional activity., tissue specificity: Abundant expression seen in the brain.,

Research Area


B Cell Receptor; Stem cell pathway; MAPK ERK Growth; MAPK G Protein; Akt PKB; Protein Acetylation

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using ATF2 (Phospho-Thr69 or 51)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838 Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from LOVO cells, using ATF2 (Phospho-Thr69 or 51) Antibody. The lane on the right is blocked with the phospho peptide.

Note

For research use only.