



Catalog #: APRab04408



Summary

CD71 (phospho Ser24) Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Host Rabbit

Application ELISA,IHC,WB, Reactivity Human, Mouse

Performance

Conjugation Unconjugated

Phospho Antibody Modification

Isotype IgG

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name TFRC

Alternative Names TFRC; Transferrin receptor protein 1; TR; TfR; TfR1; Trfr; T9; p90; CD antigen CD71

Gene ID 4155.0

P02786. The antiserum was produced against synthesized peptide derived from human **SwissProt ID**

CD71/TfR around the phosphorylation site of Ser24. AA range:15-64

Application

Dilution Ratio WB 1:500 - 1:2000 IHC 1:100 - 1:300. ELISA: 1:10000...

Molecular Weight 89kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: APRab04408




Background

This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeg, Sep 2015], function: Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the heditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site., induction: Regulated by cellular iron levels through binding of the iron regulatory proteins, IRP1 and IRP2, to iron-responsive elements in the 3'-UTR. Up-regulated upon mitogenic stimulation., miscellaneous: Canine and feline parvoviruses bind human and feline transferrin receptors and use these receptors to enter and infect cells, miscellaneous: Serum transferrin receptor (sTfR) is used as a means of detecting erythropoietin (EPO) misuse by athletes and as a diagnostic test for anemia resulting from a number of conditions including rheumatoid arthritis, pregnancy, irritable bowel syndrome and in HIV patients, PTM:N- and O-glycosylated, phosphorylated and palmitoylated. The serum form is only glycosylated., PTM: Palmitoylated on both Cys-62 and Cys-67. Cys-62 seems to be the major site of palmitoylation., PTM: Proteolytically cleaved on Arg-100 to produce the soluble serum form (sTfR), similarity: Belongs to the peptidase M28 family. M28B subfamily, similarity: Contains 1 PA (protease associated) domain, subcellular location: Identified by mass spectrometry in melanosome fractions from stage I to stage IV., subunit: Homodimer; disulfide-linked. Binds one transferrin or HFE molecule per subunit. Binds the HLA class II histocompatibility antigen, DR1. Interacts with SH3BP3. Interacts with Machupo arenavirus GPC.,

Research Area

Protein Acetylation

Image Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using CD71/TfR (Phospho-Ser24) Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab04408






Immunohistochemistry analysis of paraffin-embedded human brain, using CD71/TfR (Phospho-Ser24) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with PMA 125ng/ml 30 ', using CD71/TfR (Phospho-Ser24) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of 293 cells using Phospho-CD71 (S24) Polyclonal Antibody

Note

For research use only.