

Summary

Cdc25C (phospho Thr48) Rabbit Polyclonal Antibody
Rabbit Polyclonal Antibody
Rabbit
ELISA,IHC,WB
Human, Rat, Mouse

Performance

Conjugation	Unconjugated	
Modification	Phospho Antibody	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

Immunogen

Gene Name	CDC25C	
Alternative Names	CDC25C; M-phase inducer phosphatase 3; Dual specificity phosphatase Cdc25C	
Gene ID	995.0	
SwissProt ID	P30307.The antiserum was produced against synthesized peptide derived from human	
	CDC25C around the phosphorylation site of Thr48. AA range:14-63	

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000
Molecular Weight	80kD



Background

cell division cycle 25C(CDC25C) Homo sapiens This gene encodes a conserved protein that plays a key role in the regulation of cell division. The encoded protein directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It also suppresses p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described. [provided by RefSeq, Dec 2015],catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,developmental stage:Expressed predominantly in G2 phase.,function:Functions as a dosage-dependent inducer in mitotic control. It is a tyrosine protein phosphatase required for progression of the cell cycle. It directly dephosphorylates CDC2 and activate its kinase activity.,PTM:Phosphorylated by CHK1 on Ser-216. This phosphorylation creates a binding site for 14-3-3 protein and inhibits the phosphatase.,similarity:Belongs to the MPI phosphatase family.,similarity:Contains 1 rhodanese domain.,subunit:Interacts with HIV-1 Vpr, thereby inactivating CDC25C phosphatase activity.,

Research Area

Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Oocyte meiosis;Progesterone-mediated oocyte maturation;





Image Data


Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using CDC25C (Phospho-Thr48) Antibody



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using CDC25C (Phospho-Thr48) Antibody.



The picture on the right is blocked with the phospho peptide. $1 \quad 2$



Western blot analysis of lysates from RAW264.7 cells, using CDC25C (Phospho-Thr48) Antibody. The lane on the right is blocked with the phospho peptide.

Note

For research use only.