Polyclonal Antibody Catalog #: APRab04447



Summary

Production Name c-Fos (phospho Ser362) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application ELISA,IHC,WB, **Reactivity** Human,Mouse,Rat

Performance

Conjugation Unconjugated

Modification Phospho Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name FOS

FOS; G0S7; Proto-oncogene c-Fos; Cellular oncogene fos; G0/G1 switch regulatory Alternative Names

protein 7

Gene ID 2353.0

P01100.The antiserum was produced against synthesized peptide derived from human SwissProt ID

Fos around the phosphorylation site of Ser362. AA range:331-380

Application

Dilution Ratio WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000...

Molecular Weight 62kD

Polyclonal Antibody Catalog #: APRab04447




Background

The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death. [provided by RefSeq, Jul 2008], function: Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, c-fos and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation., PTM:Constitutively sumoylated by SUMO1, SUMO2 and SUMO3. Desumoylated by SENP2. Sumoylation requires heterodimerization with JUN and is enhanced by mitogen stimulation. Sumoylation inhibits the AP-1 transcriptional activity and is, itself, inhibited by Ras-activated phosphorylation on Thr-232.,PTM:Phosphorylated in the C-terminal upon stimulation by nerve growth factor (NGF) and epidermal growth factor (EGF). Phosphorylated, in vitro, by MAPK and RSK1. Phosphorylation on both Ser-362 and Ser-374 by MAPK1/2 and RSK1/2 leads to protein stabilization with phosphorylation on Ser-374 being the major site for protein stabilization on NGF stimulation. Phosphorylation on Ser-362 and Ser-374 primes further phosphorylations on Thr-325 and Thr-331 through promoting docking of MAPK to the DEF domain. Phosphorylation on Thr-232, induced by HA-RAS, activates the transcriptional activity and antagonizes sumoylation. Phosphorylation on Ser-362 by RSK2 in osteoblasts contributes to osteoblast transformation., similarity: Belongs to the bZIP family., similarity: Belongs to the bZIP family. Fos subfamily, similarity: Contains 1 bZIP domain, subunit: Heterodimer with JUN. Interacts with DSIPI; this interaction inhibits the binding of active AP1 to its target DNA. Interacts with MAFB.,

Research Area

MAPK ERK Growth; MAPK G Protein; Toll Like; T Cell Receptor; B Cell Antigen; Pathways in cancer; Colorectal cancer;

Image Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-


Polyclonal Antibody Catalog #: APRab04447



Phosphopeptide (Phospho-right), using Fos (Phospho-Ser362) Antibody



Immunohistochemistry analysis of paraffin-embedded human thyroid gland, using Fos (Phospho-Ser362) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from K562 cells treated with forskolin 40nM 30 ', using Fos (Phospho-Ser362) Antibody.


The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-c-Fos (S362) Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Polyclonal Antibody Catalog #: APRab04447





Western Blot analysis of 293T cells using Phospho-c-Fos (S362) Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Note

For research use only.