

Product Name: eIF2 α (phospho Ser51) Rabbit Polyclonal Antibody
Catalog #: APRab04597

Summary

Production Name	eIF2 α (phospho Ser51) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IF,ELISA
Reactivity	Human,Mouse,Rat,Dog,Pig,Fish

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	EIF2S1
Alternative Names	EIF2S1; EIF2A; Eukaryotic translation initiation factor 2 subunit 1; Eukaryotic translation initiation factor 2 subunit alpha; eIF-2-alpha; eIF-2A; eIF-2alpha
Gene ID	1965.0
SwissProt ID	P05198.The antiserum was produced against synthesized peptide derived from human eIF2 alpha around the phosphorylation site of Ser51. AA range:21-70

Application

Dilution Ratio	IF 1:50-200 WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000. Not yet tested in other applications.
-----------------------	---

Product Name: eIF2 α (phospho Ser51) Rabbit Polyclonal Antibody
Catalog #: APRab04597


Molecular Weight 38kD

Background

The translation initiation factor EIF2 catalyzes the first regulated step of protein synthesis initiation, promoting the binding of the initiator tRNA to 40S ribosomal subunits. Binding occurs as a ternary complex of methionyl-tRNA, EIF2, and GTP. EIF2 is composed of 3 nonidentical subunits, the 36-kD EIF2-alpha subunit (EIF2S1), the 38-kD EIF2-beta subunit (EIF2S2; MIM 603908), and the 52-kD EIF2-gamma subunit (EIF2S3; MIM 300161). The rate of formation of the ternary complex is modulated by the phosphorylation state of EIF2-alpha (Ernst et al., 1987 [PubMed 2948954]).[supplied by OMIM, Feb 2010],function:Functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B.,PTM:Substrate for at least 4 kinases: EIF2AK3/PERK, GCN2, HRI and PKR. Phosphorylation stabilizes the eIF-2/GDP/eIF-2B complex and prevents GDP/GTP exchange reaction, thus impairing the recycling of eIF-2 between successive rounds of initiation and leading to global inhibition of translation. In case of infection by vaccinia virus or rotavirus A, eIF2S1 phosphorylation state is modulated.,similarity:Belongs to the eIF-2-alpha family.,similarity:Contains 1 S1 motif domain.,subunit:Heterotrimer composed of an alpha, a beta and a gamma chain. Component of an EIF2 complex at least composed of CUGBP1, CALR, CALR3, EIF2S1, EIF2S2, HSP90B1 and HSPA5.,


Research Area

Image Data




Western blot analysis of lysates from K562 cells treated with IFN-alpha 1000U/ml 18h, using eIF2 alpha (Phospho-Ser51) Antibody. The lane on the right is blocked with the phospho peptide.


Product Name: eIF2 α (phospho Ser51) Rabbit Polyclonal Antibody
Catalog #: APRab04597



Immunofluorescence analysis of rat-heart tissue. 1,eIF2 α (phospho Ser51) Polyclonal Antibody (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of rat-heart tissue. 1,eIF2 α (phospho Ser51) Polyclonal Antibody (red) was diluted at 1:200 (4°C,overnight) . 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min) .3, Picture B: DAPI (blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Western Blot analysis of various cells using Phospho-eIF2 α (S51) Polyclonal Antibody diluted at 1 : 2000

Product Name: eIF2 α (phospho Ser51) Rabbit Polyclonal Antibody
Catalog #: APRab04597



Western Blot analysis of 293 cells using Phospho-eIF2 α (S51) Polyclonal Antibody diluted at 1: 2000

Note

For research use only.