

Product Name: FoxO1/3/4 (phospho Thr24/32) Rabbit Polyclonal Antibody
Catalog #: APRab04697



Summary

Production Name	FoxO1/3/4 (phospho Thr24/32) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	FOXO1/3/4 FOXO1; FKHR; FOXO1A; Forkhead box protein O1; Forkhead box protein O1A; Forkhead in rhabdomyosarcoma; FOXO3; FKHL1; FOXO3A; Forkhead box protein O3; AF6q21 protein; Forkhead in rhabdomyosarcoma-like 1; FOXO4; AFX; AFX1; MLLT7; Forkhead box
Alternative Names	
Gene ID	2308/4303 Q12778/O43524/P98177.The antiserum was produced against synthesized peptide derived from human FOXO1/3/4-pan around the phosphorylation site of Thr24/32. AA range:15-64
SwissProt ID	

Application

Product Name: FoxO1/3/4 (phospho Thr24/32) Rabbit Polyclonal Antibody
Catalog #: APRab04697



Dilution Ratio WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000..

Molecular Weight 78kD


Background

This gene belongs to the forkhead family of transcription factors which are characterized by a distinct forkhead domain. The specific function of this gene has not yet been determined; however, it may play a role in myogenic growth and differentiation. Translocation of this gene with PAX3 has been associated with alveolar rhabdomyosarcoma. [provided by RefSeq, Jul 2008],disease:Chromosomal aberrations involving FOXO1 are a cause of rhabdomyosarcoma 2 (RMS2) [MIM:268220]; also known as alveolar rhabdomyosarcoma. Translocation (2;13)(q35;q14) with PAX3; translocation t(1;13)(p36;q14) with PAX7. The resulting protein is a transcriptional activator.,function:Transcription factor.,PTM:Phosphorylated by AKT1; insulin-induced (By similarity). IGF1 rapidly induces phosphorylation of Ser-256, Thr-24, and Ser-319. Phosphorylation of Ser-256 decreases DNA-binding activity and promotes the phosphorylation of Thr-24, and Ser-319, permitting phosphorylation of Ser-322 and Ser-325, probably by CK1, leading to nuclear exclusion and loss of function. Phosphorylation of Ser-329 is independent of IGF1 and leads to reduced function. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 fork-head DNA-binding domain.,subcellular location:Shuttles between cytoplasm and nucleus.,subunit:Interacts with LRPPRC.,tissue specificity:Ubiquitous.,

Research Area


Insulin Receptor; B Cell Receptor; Protein_Acetylation

Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using FOXO1/3/4-pan (Phospho-Thr24/32) Antibody. The picture on the right is blocked with the phospho peptide.

Product Name: FoxO1/3/4 (phospho Thr24/32) Rabbit Polyclonal Antibody
Catalog #: APRab04697



Western blot analysis of lysates from 293 cells treated with Serum 20% 15', using FOXO1/3/4-PAN (Phospho-Thr24/32) Antibody. The lane on the right is blocked with the phospho peptide.

Note

For research use only.