

## Summary

Production Name	HNF4- $\alpha$ (phospho Ser313) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	ELISA,IF,IHC,WB
Reactivity	Human, Mouse, Rat

#### Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

#### Immunogen

Gene Name	HNF4A
	HNF4A; HNF4; NR2A1; TCF14; Hepatocyte nuclear factor 4-alpha; HNF-4-alpha;
Alternative Names	Nuclear receptor subfamily 2 group A member 1; Transcription factor 14; TCF-14;
	Transcription factor HNF-4
Gene ID	3172.0
SwissProt ID	P41235.The antiserum was produced against synthesized peptide derived from human
	HNF4 alpha around the phosphorylation site of Ser313. AA range:280-329

# Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in
----------------	--

## Product Name: HNF4-α (phospho Ser313) Rabbit Polyclonal Antibody Catalog #: APRab04786



other applications.

Molecular Weight

52kD

## Background


The protein encoded by this gene is a nuclear transcription factor which binds DNA as a homodimer. The encoded protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This gene may play a role in development of the liver, kidney, and intestines. Mutations in this gene have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I. Alternative splicing of this gene results in multiple transcript variants encoding several different isoforms. [provided by RefSeq, Apr 2012], alternative products: Additional isoforms seem to exist, disease: Defects in HNF4A are the cause of maturity onset diabetes of the young type 1 (MODY1) [MIM:125850]; also shortened MODY-1. MODY [MIM:606391] is a form of diabetes that is characterized by an autosomal dominant mode of inheritance, onset in childhood or early adulthood (usually before 25 years of age) and a primary defect in insulin secretion. The clinical phenotype of MODY1 is characterized by severe insulin secretory defects, and by major hyperglycemia associated with microvascular complications, function: Transcriptionally controlled transcription factor. Binds to DNA sites required for the transcription of alpha 1-antitrypsin, apolipoprotein CIII, transthyretin genes and HNF1-alpha. May be essential for development of the liver, kidney and intestine.,miscellaneous:Binds fatty acids.,online information:Hepatocyte nuclear factors entry, PTM: Phosphorylated on tyrosine residue(s); phosphorylation is important for its DNA-binding activity. Phosphorylation may directly or indirectly play a regulatory role in the subnuclear distribution, similarity: Belongs to the nuclear hormone receptor family, similarity: Belongs to the nuclear hormone receptor family. NR2 subfamily., similarity: Contains 1 nuclear receptor DNA-binding domain., subunit: Homodimerization is required for HNF4alpha to bind to its recognition site.,

#### **Research Area**


Stem cell pathway; AMPK; Protein\_Acetylation

## Image Data






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HNF4 alpha (Phospho-Ser313) Antibody




Immunofluorescence analysis of LOVO cells, using HNF4 alpha (Phospho-Ser313) Antibody. The picture on the right is blocked with the phospho peptide.




Immunohistochemistry analysis of paraffin-embedded human liver carcinoma, using HNF4 alpha (Phospho-Ser313) Antibody. The picture on the right is blocked with the phospho peptide.





Western blot analysis of lysates from HUVEC cells treated with EGF 200ng/ml 30 ', using HNF4 alpha (Phospho-Ser313) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-HNF4- $\alpha$  (S313) Polyclonal Antibody diluted at 1: 1000

#### Note

For research use only.