

Catalog #: APRab04795



## Summary

Hrs (phospho Tyr334) Rabbit Polyclonal Antibody **Production Name** 

Description Rabbit Polyclonal Antibody

Rabbit Host

**Application** ELISA, IF, IHC, WB Reactivity Human, Mouse, Rat

### **Performance**

Conjugation Unconjugated

Modification Phospho Antibody

Isotype IgG

**Clonality** Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

Gene Name HGS

HGS; HRS; Hepatocyte growth factor-regulated tyrosine kinase substrate; Hrs; Protein **Alternative Names** 

pp110

Gene ID 9146.0

O14964.The antiserum was produced against synthesized peptide derived from human SwissProt ID

HRS around the phosphorylation site of Tyr334. AA range:301-350

## **Application**

**Dilution Ratio** WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. IHC 1:100 - 1:300.

**Molecular Weight** 86kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

**Antibody** 

Catalog #: APRab04795



### **Background**

The protein encoded by this gene regulates endosomal sorting and plays a critical role in the recycling and degradation of membrane receptors. The encoded protein sorts monoubiquitinated membrane proteins into the multivesicular body, targeting these proteins for lysosome-dependent degradation. [provided by RefSeq, Dec 2010],domain:Has a double-sided UIM that can bind 2 ubiquitin molecules, one on each side of the helix, function: Involved in intracellular signal transduction mediated by cytokines and growth factors. When associated with STAM, it suppresses DNA signaling upon stimulation by IL-2 and GM-CSF. Could be a direct effector of PI3-kinase in vesicular pathway via early endosomes and may regulate trafficking to early and late endosomes by recruiting clathrin. May concentrate ubiquitinated receptors within clathrincoated regions. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with STAM (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes. May contribute to the efficient recruitment of SMADs to the activin receptor complex., PTM: Phosphorylated on Tyr-334. A minor site of phosphorylation on Tyr-329 is detected (By similarity). Phosphorylation occurs in response to EGF, IL-2, GM-CSF and HGF., similarity: Contains 1 FYVE-type zinc finger., similarity: Contains 1 UIM (ubiquitin-interacting motif) repeat., similarity: Contains 1 VHS domain., subunit: Component of the ESCRT-0 complex composed of STAM or STAM2 and HGS. Part of a complex at least composed of HSG, STAM2 (or probably STAM) and EPS15. Interacts with STAM. Interacts with STAM2. Interacts with EPS15; the interaction is direct, calcium-dependent and inhibited by SNAP25. Interacts with NF2; the interaction is direct. Interacts with ubiquitin; the interaction is direct. Interacts with VPS37C. Interacts with SMAD1, SMAD2 and SMAD3. Interacts with TSG101; the interaction mediates the association with the ESCRT-I complex. Interacts with SNAP25; the interaction is direct and decreases with addition of increasing concentrations of free calcium. Interacts with SNX1; the interaction is direct. Component of a 550 kDa membrane complex at least composed of HGS and SNX1 but excluding EGFR, tissue specificity: Ubiquitous expression in adult and fetal tissues with higher expression in testis and peripheral blood leukocytes.,

### Research Area

Endocytosis;


### **Image Data**

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab04795






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using HRS (Phospho-Tyr334) Antibody




Immunofluorescence analysis of HeLa cells treated with Forskolin 40nM 15 ', using HRS (Phospho-Tyr334) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using HRS (Phospho-Tyr334) Antibody. The picture on the right is blocked with the phospho peptide.



Catalog #: APRab04795



Western blot analysis of lysates from HepG2 cells treated with PMA 125ng/ml 30 ', using HRS (Phospho-Tyr334) Antibody. The lane on the right is blocked with the phospho peptide.

# Note

For research use only.