

# Summary

| Production Name     | IL-10R $\alpha$ (phospho Tyr496) Rabbit Polyclonal Antibody |
|---------------------|---|
| Description         | Rabbit Polyclonal Antibody                                  |
| Host                | Rabbit  |
| Application         | WB,ELISA  |
| Reactivity          | Human,Rat,Mouse   |
| Host<br>Application | Rabbit<br>WB,ELISA  |

# Performance

| Conjugation  | Unconjugated   |
|--------------|--|
| Modification | Phospho Antibody   |
| lsotype      | lgG  |
| Clonality    | Polyclonal   |
| Form         | Liquid   |
| Storage      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw   |
|              | cycles.  |
| Buffer       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification  |

# Immunogen

| Gene Name         | IL10RA  |
|-------------------|---|
| Alternative Names | IL10RA; IL10R; Interleukin-10 receptor subunit alpha; IL-10 receptor subunit alpha; IL- |
|                   | 10R subunit alpha; IL-10RA; CDw210a; Interleukin-10 receptor subunit 1; IL-10R subunit  |
|                   | 1; IL-10R1; CD antigen CD210  |
| Gene ID           | 3587.0  |
| SwissProt ID      | Q13651. The antiserum was produced against synthesized peptide derived from human       |
|                   | IL-10R alpha around the phosphorylation site of Tyr496. AA range:462-511                |

# Application

| <b>Dilution Ratio</b> | WB 1:500 - 1:2000. ELISA: 1:20000 |
|-----------------------|-----------------------------------|
| Molecular Weight      | 63kD                              |




# Background

The protein encoded by this gene is a receptor for interleukin 10. This protein is structurally related to interferon receptors. It has been shown to mediate the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines. This receptor is reported to promote survival of progenitor myeloid cells through the insulin receptor substrate-2/PI 3-kinase/AKT pathway. Activation of this receptor leads to tyrosine phosphorylation of JAK1 and TYK2 kinases. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene. [provided by RefSeq, Jan 2009],function:Receptor for IL10; binds IL10 with a high affinity.,similarity:Belongs to the type II cytokine receptor family.,tissue specificity:Spleen, thymus, and PBMC. Weak expression in pancreas, skeletal muscle, brain, heart, and kidney. Placenta, lung, and liver showed intermediate levels. Monocytes, B-cells, large granular lymphocytes, and T-cells express high levels.,

### **Research Area**


Cytokine-cytokine receptor interaction;Jak\_STAT;

### **Image Data**



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IL-10R alpha (Phospho-Tyr496) Antibody





Western blot analysis of lysates from HUVEC cells, using IL-10R alpha (Phospho-Tyr496) Antibody. The lane on the right is blocked with the phospho peptide.

### Note

For research use only.