

**Product Name: ILK (phospho Ser246) Rabbit Polyclonal Antibody**  
**Catalog #: APRab04845**



## Summary

<b>Production Name</b>	ILK (phospho Ser246) Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phospho Antibody
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	ILK
<b>Alternative Names</b>	ILK; ILK1; ILK2; Integrin-linked protein kinase; 59 kDa serine/threonine-protein kinase; ILK-1; ILK-2; p59ILK
<b>Gene ID</b>	3611.0
<b>SwissProt ID</b>	Q13418.The antiserum was produced against synthesized peptide derived from human ILK around the phosphorylation site of Ser246. AA range:212-261

## Application

<b>Dilution Ratio</b>	IHC 1:100-1:300 ELISA: 1:5000
<b>Molecular Weight</b>	

**Product Name: ILK (phospho Ser246) Rabbit Polyclonal Antibody**  
**Catalog #: APRab04845**



---


## Background

This gene encodes a protein with a kinase-like domain and four ankyrin-like repeats. The encoded protein associates at the cell membrane with the cytoplasmic domain of beta integrins, where it regulates integrin-mediated signal transduction. Activity of this protein is important in the epithelial to mesenchymal transition, and over-expression of this gene is implicated in tumor growth and metastasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2013], catalytic activity: ATP + a protein = ADP + a phosphoprotein., domain: A PH-like domain is involved in phosphatidylinositol phosphate binding., enzyme regulation: Stimulated rapidly but transiently by both cell fibronectin interactions, as well as by insulin, in a PI3-K-dependent manner, likely via the binding of PtdIns(3,4,5)P3 with a PH-like domain of ILK., function: Receptor-proximal protein kinase regulating integrin-mediated signal transduction. May act as a mediator of inside-out integrin signaling. Focal adhesion protein part of the complex ILK-PINCH. This complex is considered to be one of the convergence points of integrin- and growth factor-signaling pathway. Could be implicated in mediating cell architecture, adhesion to integrin substrates and anchorage-dependent growth in epithelial cells. Phosphorylates beta-1 and beta-3 integrin subunit on serine and threonine residues, but also AKT1 and GSK3B., PTM: Autophosphorylated on serine residues., similarity: Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family., similarity: Contains 1 protein kinase domain., similarity: Contains 5 ANK repeats., subunit: Interacts with cytoplasmic domain of beta 1 subunit of integrin. Could also interact with beta 2, beta 3 and/or beta 5 subunit of integrin. Interacts (via ANK repeats) with LIMS1 and LIMS2. Interacts with parvins and probably TGFBI1., tissue specificity: Highly expressed in heart followed by skeletal muscle, pancreas and kidney. Weakly expressed in placenta, lung and liver.,

## Research Area

PPAR; Focal adhesion; Endometrial cancer;

## Image Data



Immunohistochemistry analysis of paraffin-embedded human brain, using ILK (Phospho-Ser246) Antibody. The picture on the right is blocked with the phospho peptide.

**Product Name: ILK (phospho Ser246) Rabbit Polyclonal  
Antibody  
Catalog #: APRab04845**

---



**Note**

For research use only.