# Product Name: IRAK-1 (phospho Thr387) Rabbit

Polyclonal Antibody Catalog #: APRab04867



## **Summary**

**Production Name** IRAK-1 (phospho Thr387) Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application Elisa

**Reactivity** Human, Mouse, Rat

#### **Performance**

| Conjugation  | Unconjugated   |
|--------------|--|
| Modification | Phospho Antibody   |
| Isotype      | IgG  |
| Clonality    | Polyclonal   |
| Form         | Liquid   |
| Storage      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.       |
| Purification | Affinity purification  |

### **Immunogen**

Gene Name IRAK1

Alternative Names IRAK1; IRAK; Interleukin-1 receptor-associated kinase 1; IRAK-1

Gene ID 3654.0

P51617.The antiserum was produced against synthesized peptide derived from human

IRAK1 around the phosphorylation site of Thr387. AA range:353-402

## **Application**

SwissProt ID

**Dilution Ratio** ELISA: 1:10000

**Molecular Weight** 

### **Background**

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# Product Name: IRAK-1 (phospho Thr387) Rabbit

Polyclonal Antibody Catalog #: APRab04867




This gene encodes the interleukin-1 receptor-associated kinase 1, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. This gene is partially responsible for IL1-induced upregulation of the transcription factor NF-kappa B. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, function:Binds to the IL-1 type I receptor following IL-1 engagement, triggering intracellular signaling cascades leading to transcriptional up-regulation and mRNA stabilization. Isoform 1 binds rapidly but is then degraded allowing isoform 2 to mediate a slower, more sustained response to the cytokine. Isoform 2 is inactive suggesting that the kinase activity of this enzyme is not required for IL-1 signaling. Once phosphorylated, IRAK1 recruits the adapter protein PELI1,PTM:Autophosphorylated or is transphosphorylated by IRAK4 following recruitment to the IL-1RI. In the case of isoform 1, this is linked to ubiquitination and degradation, similarity:Belongs to the protein kinase superfamily, similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Pelle subfamily, similarity:Contains 1 protein kinase domain, subunit:IL-1 stimulation leads to the formation of a signaling complex which dissociates from the IL-1 receptor following the binding of PELI1. Interacts with IL1RL1. Interacts with IRAK1BP1., tissue specificity:Isoform 1 and isoform 2 are ubiquitously expressed in all tissues examined, with isoform 1 being more strongly expressed than isoform 2.

#### Research Area

Apoptosis Inhibition; Apoptosis Mitochondrial; Apoptosis Overview; Toll Like; Neurotrophin;

### **Image Data**



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using IRAK1 (Phospho-Thr387) Antibody

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838