

## Summary

Synapsin I (phospho Ser9) Rabbit Polyclonal Antibody
Rabbit Polyclonal Antibody
Rabbit
WB,
Human, Mouse, Rat

#### Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

#### Immunogen

Gene Name	SYN1
Alternative Names	SYN1; Synapsin-1; Brain protein 4.1; Synapsin I
Gene ID	6853.0
SwissProt ID	P17600.The antiserum was produced against synthesized peptide derived from human
	Synapsin around the phosphorylation site of Ser9. AA range:3-52

# Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.
Molecular Weight	77kD




#### Background

This gene is a member of the synapsin gene family. Synapsins encode neuronal phosphoproteins which associate with the cytoplasmic surface of synaptic vesicles. Family members are characterized by common protein domains, and they are implicated in synaptogenesis and the modulation of neurotransmitter release, suggesting a potential role in several neuropsychiatric diseases. This member of the synapsin family plays a role in regulation of axonogenesis and synaptogenesis. The protein encoded serves as a substrate for several different protein kinases and phosphorylation may function in the regulation of this protein in the nerve terminal. Mutations in this gene may be associated with X-linked disorders with primary neuronal degeneration such as Rett syndrome. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008], disease: Defects in SYN1 are a cause of epilepsy Xlinked with variable learning disabilities and behavior disorders [MIM:300491]. XELBD is characterized by variable combinations of epilepsy, learning difficulties, macrocephaly, and aggressive behavior., function: Neuronal phosphoprotein that coats synaptic vesicles, binds to the cytoskeleton, and is believed to function in the regulation of neurotransmitter release. The complex formed with NOS1 and CAPON proteins is necessary for specific nitric-oxid functions at a presynaptic level, PTM: Substrate of at least four different protein kinases. It is probable that phosphorylation plays a role in the regulation of synapsin-1 in the nerve terminal. Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the synapsin family., subunit: Homodimer. Interacts with CAPON. Forms a ternary complex with NOS1. Isoform Ib interacts with PRNP.,


### **Research Area**

#### Image Data




Western blot analysis of lysates from 293 cells treated with PMA 200nM 30 ', using Synapsin (Phospho-Ser9) Antibody. The lane on the right is blocked with the phospho peptide.






Western Blot analysis of various cells using Phospho-Synapsin I (S9) Polyclonal Antibody diluted at 1: 1000



Western Blot analysis of 3T3 cells using Phospho-Synapsin I (S9) Polyclonal Antibody diluted at 1: 1000



Western blot analysis of HELA KB 293T 3T3 lysis using Phospho-Synapsin I (S9) antibody. Antibody was diluted at 1:1000

#### Note

For research use only.