

**Product Name: MER/TYRO3 (Phospho-Tyr753/Tyr685)  
Rabbit Polyclonal Antibody  
Catalog #: APRab05800**



## Summary

<b>Production Name</b>	MER/TYRO3 (Phospho-Tyr753/Tyr685) Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human:Y753/Y685,Mouse:Y748/Y675,Rat:Y748/Y675

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phospho Antibody
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	PBS, pH 7.4, containing 0.02% New type preservative N as Preservative and 50% Glycerol.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	MERTK MER
<b>Alternative Names</b>	MERTK MER
<b>Gene ID</b>	10461/7301
<b>SwissProt ID</b>	Q12866/Q06418.Synthesized phospho-peptide around the phosphorylation site of human MER/TYRO3 (Phospho-Tyr753/Tyr685)

## Application

<b>Dilution Ratio</b>	WB 1:500-2000;ELISA 2000-20000
<b>Molecular Weight</b>	110kD

**Product Name: MER/TYRO3 (Phospho-Tyr753/Tyr685)  
Rabbit Polyclonal Antibody  
Catalog #: APRab05800**




## Background

This gene is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in this gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP). [provided by RefSeq, Jul 2008],catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in MERTK are a cause of retinitis pigmentosa (RP) [MIM:268000]. RP that leads to degeneration of retinal photoreceptor cells. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well.,function:In case of filovirus infection, seems to function as a cell entry factor.,online information:Retina International's Scientific Newsletter,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. AXL/UFO subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 fibronectin type-III domains.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines.,

## Research Area

## Image Data



Western Blot analysis of 293T cells using Antibody diluted at 500. Secondary antibody was diluted at 1:20000

## Note

For research use only.