

Product Name: ZAK (Phospho-Ser165) Rabbit Polyclonal Antibody
Catalog #: APRab06118

Summary

Production Name	ZAK (Phospho-Ser165) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MLTK ZAK HCCS4 Mitogen-activated protein kinase kinase kinase MLT (EC 2.7.11.25) (Human cervical cancer suppressor gene 4 protein) (HCCS-4) (Leucine zipper- and sterile alpha motif-containing kinase) (MLK-like mitogen-activated protein triple kinase) (Mixed lineage kinase-related kinase) (MLK-related kinase) (MRK) (Sterile alpha motif- and leucine zipper-containing kinase AZK)
Alternative Names	
Gene ID	51776.0
SwissProt ID	Q9NYL2.

Application

Dilution Ratio	WB 1:500-2000
Molecular Weight	88kD


Product Name: ZAK (Phospho-Ser165) Rabbit Polyclonal Antibody
Catalog #: APRab06118

Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation by PKN1 and autophosphorylation on Thr-161 and Ser-165.,function:Stress-activated component of a protein kinase signal transduction cascade. Regulates the JNK and p38 pathways. Pro-apoptotic. Role in regulation of S and G2 cell cycle checkpoint by direct phosphorylation of CHEK2. Isoform 1, but not isoform 2, causes cell shrinkage and disruption of actin stress fibers. Isoform 1 may have role in neoplastic cell transformation and cancer development. Isoform 1, but not isoform 2, phosphorylates histone H3 at 'Ser-28'.similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,subcellular location:Translocates to the nucleus upon ultraviolet B irradiation.,subunit:Homodimer. Interacts with PKN1 and ZNF33A.,tissue specificity:Ubiquitously expressed. Isoform 2 is the predominant form in all tissues examined, except for liver, in which isoform 1 is more highly expressed.,

Research Area

Image Data



Western Blot analysis of HL-60 cell ,using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

Note

For research use only.