# **Product Name: ABCF1 Rabbit Polyclonal Antibody**

Catalog #: APRab06420



# **Summary**

Production Name ABCF1 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB

**Reactivity** Human, Mouse, Rat

# **Performance**

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

### **Immunogen**

Gene Name ABCF1

ABCF1; ABC50; ATP-binding cassette sub-family F member 1; ATP-binding cassette 50;

Alternative Names

TNF-alpha-stimulated ABC protein

Gene ID 23.0

SwissProt ID Q8NE71.Synthesized peptide derived from ABCF1 . at AA range: 760-840

## **Application**

**Dilution Ratio** WB 1:500-1:2000. ELISA: 1:10000.

Molecular Weight 95kD

# **Background**

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins

# **Product Name: ABCF1 Rabbit Polyclonal Antibody**


Catalog #: APRab06420



transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the GCN20 subfamily. Unlike other members of the superfamily, this protein lacks the transmembrane domains which are characteristic of most ABC transporters. This protein may be regulated by tumor necrosis factor-alpha and play a role in enhancement of protein synthesis and the inflammation process. [provided by RefSeq, Jul 2008],function:May function in mRNA translation.,induction:By TNF (in cultured synoviocytes).,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the ABC transporter family.,similarity:Belongs to the ABC transporter family. EF3 subfamily,,similarity:Contains 2 ABC transporter domains.,subunit:Interacts with eukaryotic initiation factor 2 (eIF2). Associates with the ribosomes,,tissue specificity:Ubiquitous.,

### Research Area

## **Image Data**



Western Blot analysis of HeLa cells using ABCF1 Polyclonal Antibody

### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838