

# Summary

Production Name	ABHD12 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ELISA
Reactivity	Human, Mouse, Rat

### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at $4^{\circ}$ C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

## Immunogen

Gene Name	ABHD12	
Alternative Names	ABHD12; C20orf22; Monoacylglycerol lipase ABHD12; 2-arachidonoylglycerol	
	hydrolase; Abhydrolase domain-containing protein 12	
Gene ID	26090.0	
SwissProt ID	Q8N2K0.The antiserum was produced against synthesized peptide derived from	
	human ABHD12. AA range:231-280	

# Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000
Molecular Weight	45kD




### Background


This gene encodes an enzyme that catalyzes the hydrolysis of 2-arachidonoyl glycerol (2-AG), the main endocannabinoid lipid transmitter that acts on cannabinoid receptors, CB1 and CB2. The endocannabinoid system is involved in a wide range of physiological processes, including neurotransmission, mood, appetite, pain appreciation, addiction behavior, and inflammation. Mutations in this gene are associated with the neurodegenerative disease, PHARC (polyneuropathy, hearing loss, ataxia, retinitis pigmentosa, and cataract), resulting from an inborn error of endocannabinoid metabolism. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.[provided by RefSeq, Jan 2011],catalytic activity:Hydrolyzes glycerol monoesters of long-chain fatty acids.,function:Has 2-arachidonoylglycerol hydrolase activity (By similarity). May be a regulator of endocannabinoid signaling pathways.,PTM:Glycosylated.,similarity:Belongs to the serine esterase family.,


### **Research Area**

Image Data




Western blot analysis of lysates from COLO cells, using ABHD12 Antibody. The lane on the right is blocked with the









Immunohistochemical analysis of paraffin-embedded Human ovary. Antibody was diluted at 1:100 (4°,overnight) . Highpressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

**Note** For research use only.