## **Product Name: ATP5A Rabbit Polyclonal Antibody**

Catalog #: APRab07324



### **Summary**

Production Name ATP5A Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB,

**Reactivity** Human, Mouse, Rat

### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

Storage

Gene Name ATP5A1

Alternative Names ATP5A1; ATP5A; ATP5AL2; ATPM; ATP synthase subunit alpha; mitochondrial

**Gene ID** 498.0

P25705. The antiserum was produced against synthesized peptide derived from human

ATP5A1. AA range:201-250

## **Application**

**SwissProt ID** 

**Dilution Ratio** WB 1:500 - 1:2000. ELISA: 1:40000...

Molecular Weight 60kD

## **Background**

# Product Name: ATP5A Rabbit Polyclonal Antibody Catalog #: APRab07324




This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, using an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel consists of three main subunits (a, b, c). This gene encodes the alpha subunit of the catalytic core. Alternatively spliced transcript variants encoding the different isoforms have been identified. Pseudogenes of thifunction: Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Subunits alpha and beta form the catalytic core in F(1). Rotation of the central stalk against the surrounding alpha(3) beta(3) subunits leads to hydrolysis of ATP in three separate catalytic sites on the beta subunits. Subunit alpha does not bear the catalytic high-affinity ATP-binding sites., PTM:The N-terminus is blocked., similarity:Belongs to the ATPase alpha/beta chains family,,subcellular location:Peripheral membrane protein.,subunit:F-type ATPases have 2 components, CF(1) - the catalytic core - and CF(0) - the membrane proton channel. CF(1) has five subunits: alpha(3), beta(3), gamma(1), delta(1), epsilon(1). CF(0) has three main subunits: a, b and c. Interacts with ATPAF2.,tissue specificity:Fetal lung, heart, liver, gut and kidney. Expressed at higher levels in the fetal brain, retina and spinal cord.,

#### Research Area

Oxidative phosphorylation; Alzheimer's disease; Parkinson's disease; Huntington's disease;


### **Image Data**




Western blot analysis of lysates from 293 and RAW264.7 cells, using ATP5A1 Antibody. The lane on the right is blocked with the synthesized peptide.

## **Product Name: ATP5A Rabbit Polyclonal Antibody** Catalog #: APRab07324






Western blot analysis of the lysates from HT-29 cells using ATP5A1 antibody.



Western Blot analysis of various cells using ATP5A Polyclonal Antibody diluted at 1: 500



Western Blot analysis of RAW264.7 cells using ATP5A Polyclonal Antibody diluted at 1: 500

### Note

For research use only.