# **Product Name: AVP Receptor V2 Rabbit Polyclonal**

**Antibody** 

Catalog #: APRab07378



## **Summary**

Production Name AVP Receptor V2 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

Application IF,WB,ELISA

**Reactivity** Human, Rat, Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw  $\bf Storage$ 

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

Gene Name AVPR2

AVPR2; ADHR; DIR3; V2R; Vasopressin V2 receptor; V2R; AVPR V2; Antidiuretic Alternative Names

hormone receptor; Renal-type arginine vasopressin receptor

**Gene ID** 554.0

P30518.The antiserum was produced against synthesized peptide derived from human SwissProt ID

AVPR2. AA range:72-121

# **Application**

**Dilution Ratio** 

WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

application

applications.

# **Product Name: AVP Receptor V2 Rabbit Polyclonal**

Antibody

Catalog #: APRab07378



Molecular Weight 38kD


## **Background**

This gene encodes the vasopressin receptor, type 2, also known as the V2 receptor, which belongs to the seventransmembrane-domain G protein-coupled receptor (GPCR) superfamily, and couples to Gs thus stimulating adenylate cyclase. The subfamily that includes the V2 receptor, the V1a and V1b vasopressin receptors, the oxytocin receptor, and isotocin and mesotocin receptors in non-mammals, is well conserved, though several members signal via other G proteins. All bind similar cyclic nonapeptide hormones. The V2 receptor is expressed in the kidney tubule, predominantly in the distal convoluted tubule and collecting ducts, where its primary property is to respond to the pituitary hormone arginine vasopressin (AVP) by stimulating mechanisms that concentrate the urine and maintain water homeostasis in the organism. When the function of this gene is lost, the disease Nephrogenic Diabetes Insipidus disease: Defects in AVPR2 are the cause of diabetes insipidus nephrogenic X-linked (XNDI) [MIM:304800]; also known as diabetes insipidus nephrogenic type 1. XNDI is caused by the inability of the renal collecting ducts to absorb water in response to arginine vasopressin. It is characterized by excessive water drinking (polydypsia), excessive urine excretion (polyuria), persistent hypotonic urine, and hypokalemia, disease: Defects in AVPR2 are the cause of nephrogenic syndrome of inappropriate antidiuresis (NSIAD) [MIM:300539]. This disorder is characterized by an inability to excrete a free water load, with inappropriately concentrated urine and resultant hyponatremia, hypoosmolarity, and natriuresis, function: Receptor for arginine vasopressin. The activity of this receptor is mediated by G proteins which activate adenylate cyclase, online information: AVPR2 pages, similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Kidney.,

#### Research Area

Neuroactive ligand-receptor interaction;

#### **Image Data**



Immunofluorescence analysis of MCF7 cells, using AVPR2 Antibody. The picture on the right is blocked with the synthesized peptide.


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: AVP Receptor V2 Rabbit Polyclonal**


**Antibody** 

Catalog #: APRab07378





Western blot analysis of lysates from RAW264.7 cells, using AVPR2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from Jurkat cells using AVPR2 antibody.

#### Note

For research use only.