

Summary

Production Name	CARP Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ANKRD1
	ANKRD1; C193; CARP; HA1A2; Ankyrin repeat domain-containing protein 1; Cardiac
Alternative Names	ankyrin repeat protein; Cytokine-inducible gene C-193 protein; Cytokine-inducible
	nuclear protein
Gene ID	27063.0
SwissProt ID	Q15327.The antiserum was produced against synthesized peptide derived from human
	ANKRD1. AA range:231-280

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000
Molecular Weight	36kD




Background

ankyrin repeat domain 1(ANKRD1) Homo sapiens The protein encoded by this gene is localized to the nucleus of endothelial cells and is induced by IL-1 and TNF-alpha stimulation. Studies in rat cardiomyocytes suggest that this gene functions as a transcription factor. Interactions between this protein and the sarcomeric proteins myopalladin and titin suggest that it may also be involved in the myofibrillar stretch-sensor system. [provided by RefSeq, Jul 2008], disease: Defects in ANKRD1 may be a cause of total anomalous pulmonary venous return (TAPVR) [MIM:106700]. TAPVR is a rare congenital heart disease (CHD) in which the pulmonary veins fail to connect to the left atrium during cardiac development, draining instead into either the right atrium or one of its venous tributaries. This disease accounts for 1.5% of all CHDs and has a prevalence of approximately 1 out of 15'000 live births., function: May play an important role in endothelial cell activation. May act as a nuclear transcription factor that negatively regulates the expression of cardiac genes. Induction seems to be correlated with apoptotic cell death in hepatoma cells.,induction:By TNF, IL1A and parthenolide, miscellaneous: A chromosomal aberration in the region of ANKRD1 may be a cause of total anomalous pulmonary venous return (TAPVR) [MIM:106700]. Translocation t(10;21)(g23.31;g11.2). The translocation apparently alteres the expression pattern of nearby genes on chromosome 10 by means of a positional effect, and among the genes whose expression pattern is changed due to this chromosomal rearrangement, the ANKRD1 stood out as a plausible candidate gene for TAPVR pathogenesis. This rearrangement apparently does not disrupts any known genes, similarity: Contains 5 ANK repeats., subunit: Interacts with YBX1 (By similarity). Interacts with TTN/titin., tissue specificity: Mainly expressed in activated vascular endothelial cells. To a lower extent, also expressed in hepatoma cells.,

Research Area


Image Data



Western blot analysis of lysates from COLO205 cells, using ANKRD1 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: CARP Rabbit Polyclonal Antibody Catalog #: APRab07933





Western Blot analysis of various cells using CARP Polyclonal Antibody

Note

For research use only.