

Summary

Production Name	CD148 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC,IF,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PTPRJ
Alternative Names	PTPRJ; DEP1; Receptor-type tyrosine-protein phosphatase eta; Protein-tyrosine phosphatase eta; R-PTP-eta; Density-enhanced phosphatase 1; DEP-1; HPTP eta; Protein-tyrosine phosphatase receptor type J; R-PTP-J; CD148
Gene ID	5795.0
SwissProt ID	Q12913.The antiserum was produced against synthesized peptide derived from the Internal region of human PTPRJ. AA range:861-910

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC-p: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200
Molecular Weight	150kD


Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes, including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region containing five fibronectin type III repeats, a single transmembrane region, and a single intracytoplasmic catalytic domain, and thus represents a receptor-type PTP. This protein is present in all hematopoietic lineages, and was shown to negatively regulate T cell receptor signaling possibly through interfering with the phosphorylation of Phospholipase C Gamma 1 and Linker for Activation of T Cells. This protein can also dephosphorylate the PDGF beta receptor, and may be involved in UV-induced signal transduction. Multiple transcript variants encoding different isoforms catalytic activity: Protein tyrosine phosphate + H₂O = protein tyrosine + phosphate., disease: Defects in PTPRJ are found in cancers of colon, lung, and breast., function: May contribute to the mechanism of contact inhibition of cell growth., PTM: N- and O-glycosylated., similarity: Belongs to the protein-tyrosine phosphatase family. Receptor class 3 subfamily., similarity: Contains 1 tyrosine-protein phosphatase domain., similarity: Contains 9 fibronectin type-III domains.,

Research Area


Adherens_Junction;

Image Data




Western blot analysis of lysate from 293, HeLa cells, using PTPRJ Antibody.


Product Name: CD148 Rabbit Polyclonal Antibody
Catalog #: APRab08216



Western Blot analysis of HeLa, 293 cells using CD148 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA) .




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100




Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100

Product Name: CD148 Rabbit Polyclonal Antibody
Catalog #: APRab08216



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100

Note

For research use only.