

## Summary

<b>Production Name</b>	CD218b Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	IL18RAP IL1R7 Interleukin-18 receptor accessory protein (IL-18 receptor accessory protein;IL-18RAcP;Accessory protein-like;AcPL;CD218 antigen-like family member B;CDw218b;IL-
<b>Alternative Names</b>	1R accessory protein-like;IL-1RAcPL;Interleukin-1 receptor 7;IL-1R-7;IL-1R7;Interleukin-18 receptor accessory protein-like;Interleukin-18 receptor beta;IL-18R-beta;IL-18Rbeta;CD antigen CD218b)
<b>Gene ID</b>	8807.0
<b>SwissProt ID</b>	O95256.Synthetic peptide from human protein at AA range: 291-340

## Application

<b>Dilution Ratio</b>	IHC-p 1:50-200, ELISA 1:10000-20000.
-----------------------	--------------------------------------

---

## Molecular Weight


## Background

The protein encoded by this gene is an accessory subunit of the heterodimeric receptor for interleukin 18 (IL18), a proinflammatory cytokine involved in inducing cell-mediated immunity. This protein enhances the IL18-binding activity of the IL18 receptor and plays a role in signaling by IL18. Mutations in this gene are associated with Crohn's disease and inflammatory bowel disease, and susceptibility to celiac disease and leprosy. Alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known. [provided by RefSeq, Feb 2014],function:Required for the high affinity binding of interleukin 18 (IL-18) to its receptor complex (By similarity). Together with IL18R1 mediates IL-18-dependent activation of NF-kappa-B and JNK.,induction:Induced by IFN-alpha and IL-12 in natural killer (NK) cells and T-cells.,similarity:Belongs to the interleukin-1 receptor family.,similarity:Contains 1 TIR domain.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Together with IL18R1, probably belongs to the IL-18 receptor complex. Seems not to interact directly with IL-18.,tissue specificity:Strongly expressed in peripheral blood leukocytes, spleen, lung, and, to a lesser extent, in colon, but not in any other tissue tested. Strongly expressed in T-cells polarized with IL-12. Specifically coexpressed with IL18R1 in Th1 cells.,

## Research Area


Cytokine-cytokine receptor interaction;

## Image Data




Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200

**Product Name: CD218b Rabbit Polyclonal Antibody**  
**Catalog #: APRab08283**



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

### **Note**

For research use only.