Product Name: CD28 Rabbit Polyclonal Antibody

Catalog #: APRab08324



Summary

Production Name CD28 Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit
Application WB

Reactivity Human, Rat, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name CD28

Alternative Names CD28; T-cell-specific surface glycoprotein CD28; TP44; CD28

Gene ID 940.0

P10747.Synthesized peptide derived from T-cell-specific surface glycoprotein CD28 at SwissProt ID

AA range: 11-60

Application

Dilution Ratio WB 1:500-1:2000. ELISA: 1:20000.

Molecular Weight 25kD

Background

The protein encoded by this gene is essential for T-cell proliferation and survival, cytokine production, and T-helper type-2

Product Name: CD28 Rabbit Polyclonal Antibody

C EnkiLife


Catalog #: APRab08324

development. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2011], function: Involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival., online information: CD28 entry, similarity: Contains 1 Ig-like V-type (immunoglobulin-like) domain., subunit: Homodimer; disulfide-linked. Interacts with DUSP14. Binds to CD80/B7-1 and CD86/B7-2/B70., tissue specificity: Expressed in T-cells and plasma cells, but not in less mature B-cells.,

Research Area

Cell adhesion molecules (CAMs);T_Cell_Receptor;Intestinal immune network for IgA production;Type I diabetes mellitus;Autoimmune thyroid disease;Systemic lupus erythematosus;Allograft rejection;Graft-versus-host disease;Viral myocarditis;

Image Data



Western Blot analysis of K562 cells using CD28 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Note

For research use only.