

## Summary

<b>Production Name</b>	CD89 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	FCAR
<b>Alternative Names</b>	FCAR; CD89; Immunoglobulin alpha Fc receptor; IgA Fc receptor; CD89
<b>Gene ID</b>	2204.0
<b>SwissProt ID</b>	P24071.The antiserum was produced against synthesized peptide derived from the Internal region of human FCAR. AA range:31-80

## Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. IHC-p: 1:100-1:300. ELISA: 1:10000..
<b>Molecular Weight</b>	32kD

## Background


**Product Name: CD89 Rabbit Polyclonal Antibody**  
**Catalog #: APRab08480**




This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008], alternative products: Additional isoforms seem to exist, function: Binds to the Fc region of immunoglobulins alpha. Mediates several functions including cytokine production., similarity: Contains 2 Ig-like C2-type (immunoglobulin-like) domains., subunit: Associates with the Fc epsilon RI gamma 2 receptor inducing tyrosine phosphorylation of gamma 2., tissue specificity: Differentially expressed between blood and mucosal myeloid cells. Monocytes express isoform A.1, isoform A.2 and isoform A.3 while alveolar macrophages express isoform A.1 and isoform A.2; however they express only one isoform at their surfaces.,

## Research Area

## Image Data



Western Blot analysis of NIH-3T3 cells using CD89 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-tonsils, antibody was diluted at 1:100

**Product Name: CD89 Rabbit Polyclonal Antibody**  
**Catalog #: APRab08480**

---



**Note**

For research use only.