Product Name: CHP Rabbit Polyclonal Antibody

Catalog #: APRab08778



Summary

Production Name CHP Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit
Application WB

Reactivity Human, Mouse, Rat

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

Gene Name CHP1

CHP1; CHP; Calcineurin B homologous protein 1; Calcineurin B-like protein; Calcium-

Alternative Names binding protein CHP; Calcium-binding protein p22; EF-hand calcium-binding domain-

containing protein p22

Gene ID 11261.0

SwissProt ID Q99653.Synthesized peptide derived from the Internal region of human CHP.

Application

Dilution Ratio WB 1:500-1:2000. ELISA: 1:20000.

Molecular Weight 24kD

Background

Product Name: CHP Rabbit Polyclonal Antibody Catalog #: APRab08778




This gene encodes a phosphoprotein that binds to the Na+/H+ exchanger NHE1. This protein serves as an essential cofactor which supports the physiological activity of NHE family members and may play a role in the mitogenic regulation of NHE1. The protein shares similarity with calcineurin B and calmodulin and it is also known to be an endogenous inhibitor of calcineurin activity. [provided by RefSeq, Jul 2008],function:Required for constitutive membrane traffic. Inhibits GTPase-stimulated Na(+)/H(+) exchange. Also inhibits calcineurin phosphatase activity. Required for activity of SLC9A1/NHE1,PTM:Both N-myristoylation and calcium-mediated conformational changes are essential for its function in exocytic traffic.,PTM:Phosphorylated; decreased phosphorylation is associated with an increase in exchange activity. The phosphorylation state may regulate the binding to NHE1, similarity:Contains 4 EF-hand domains, subunit:Monomer (By similarity). Specifically binds to SLC9A1/NHE1 at a domain that is critical for growth factor stimulation of exchange activity, tissue specificity:Ubiquitously expressed. Has been found in fetal eye, lung, liver, muscle, heart, kidney, thymus and spleen,

Research Area

MAPK ERK Growth; MAPK G Protein; Calcium; Oocyte

meiosis; Apoptosis_Inhibition; Apoptosis_Mitochondrial; Apoptosis_Overview; WNT; WNT-T CELLAxon guidance; VEGF; Natural killer cell mediated cytotoxicity; T_Cell_Receptor; B_Cell_Antigen; Long-term potentiation; Alzheimer's disease; Amyotrophic lateral sclerosis (ALS);

Image Data



Western Blot analysis of HeLa cells using CHP Polyclonal Antibody

Note

For research use only.