

Product Name: Cleaved-Factor B Bb (K260) Rabbit Polyclonal Antibody
Catalog #: APRab08986

Summary

Production Name	Cleaved-Factor B Bb (K260) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	CFB
Alternative Names	CFB; BF; BFD; Complement factor B; C3/C5 convertase; Glycine-rich beta glycoprotein; GBG; PBF2; Properdin factor B
Gene ID	629.0
SwissProt ID	P00751.The antiserum was produced against synthesized peptide derived from human CFAB Bb. AA range:241-290

Application

Dilution Ratio	WB 1:500-1:2000. ELISA: 1:20000.
Molecular Weight	57+85kD

Product Name: Cleaved-Factor B Bb (K260) Rabbit Polyclonal Antibody
Catalog #: APRab08986


Background

This gene encodes complement factor B, a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene for complement catalytic activity: Cleavage of Arg-Ser bond in complement component C3 alpha-chain to yield C3a and C3b, and Arg-Xaa bond in complement component C5 alpha-chain to yield C5a and C5b, function: Factor B which is part of the alternate pathway of the complement system is cleaved by factor D into 2 fragments: Ba and Bb. Bb, a serine protease, then combines with complement factor 3b to generate the C3 or C5 convertase. It has also been implicated in proliferation and differentiation of preactivated B-lymphocytes, rapid spreading of peripheral blood monocytes, stimulation of lymphocyte blastogenesis and lysis of erythrocytes. Ba inhibits the proliferation of preactivated B-lymphocytes, polymorphism: Two major variants, F and S, and 2 minor variants, as well as at least 14 very rare variants, have been identified. The variants His-9 and Gln-32 are associated with a reduced risk of age-related macular degeneration (ARMD) [MIM:603075]. ARMD is a multifactorial eye disease and the most common cause of irreversible vision loss in the developed world, similarity: Belongs to the peptidase S1 family, similarity: Contains 1 peptidase S1 domain, similarity: Contains 1 VWFA domain, similarity: Contains 3 Sushi (CCP/SCR) domains, subunit: Monomer,

Research Area

Complement and coagulation cascades;

Image Data



Western blot analysis of lysates from K562 cells, treated with etoposide 25uM 1h, using CFAB Bb (Cleaved-Lys260) Antibody. The lane on the right is blocked with the synthesized peptide.

**Product Name: Cleaved-Factor B Bb (K260) Rabbit
Polyclonal Antibody
Catalog #: APRab08986**



Note

For research use only.