Catalog #: APRab09235



Summary

Connexin 47 Rabbit Polyclonal Antibody
Rabbit Polyclonal Antibody
Rabbit
IF,WB,ELISA
Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storogo	Store at 4° C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
Storage	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	GJC2
Alternative Names	GJC2; GJA12; Gap junction gamma-2 protein; Connexin-46.6; Cx46.6; Connexin-47;
	Cx47; Gap junction alpha-12 protein
Gene ID	57165.0
SwissProt ID	Q5T442.The antiserum was produced against synthesized peptide derived from human
	CXG2. AA range:21-70

Application


Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.
Molecular Weight	47kD

Background

This gene encodes a gap junction protein. Gap junction proteins are members of a large family of homologous connexins and comprise 4 transmembrane, 2 extracellular, and 3 cytoplasmic domains. This gene plays a key role in central myelination and is involved in peripheral myelination in humans. Defects in this gene are the cause of autosomal recessive Pelizaeus-Merzbacher-like disease-1. [provided by RefSeq, Jul 2008],caution:It is uncertain whether Met-1 or Met-4 is the initiator.,disease:Defects in GJC2 are the cause of Leukodystrophy hypomyelinating type 2 (HLD2) [MIM:608804]; also known as Pelizaeus-Merzbacher-like disease autosomal recessive type 1. HLD2 is an autosomal recessive hypomyelinating leukodystrophy characterized by nystagmus, impaired motor development, ataxia, choreoathetotic movements, dysarthria and progressive spasticity.,function:One gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a role in myelination in central and peripheral nervous systems.,similarity:Belongs to the connexin family. Gamma-type subfamily.,subunit:A connexon is composed of a hexamer of connexins. Interacts with TJP1.,tissue specificity:Expressed in central nervous system, in sciatic nerve and sural nerve. Also detected in skeletal muscles.,


Research Area

Image Data




Immunofluorescence analysis of A549 cells, using CXG2 Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: Connexin 47 Rabbit Polyclonal Antibody **Control Control Control**



Western blot analysis of lysates from A549 cells, using CXG2 Antibody. The lane on the right is blocked with the synthesized





Note

For research use only.