

**Product Name: CRLF3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab09408**



## Summary

<b>Production Name</b>	CRLF3 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	CRLF3 CREME9 CYTOR4 P48
<b>Alternative Names</b>	
<b>Gene ID</b>	51379.0
<b>SwissProt ID</b>	Q8IUI8.Synthesized peptide derived from human protein . at AA range: 270-350

## Application

<b>Dilution Ratio</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Molecular Weight</b>	48kD

## Background

This gene encodes a cytokine receptor-like factor that may negatively regulate cell cycle progression at the G0/G1 phase. Studies of the related rat protein suggest that it may regulate neuronal morphology and synaptic vesicle biogenesis. This


**Product Name: CRLF3 Rabbit Polyclonal Antibody**  
**Catalog #: APRab09408**



gene is one of several genes located in the neurofibromatosis type I tumor suppressor region on the q arm of chromosome 17, a region that is subject to microdeletions, duplications, chromosomal breaks and rearrangements. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 2 and 5. [provided by RefSeq, Aug 2012],similarity:Belongs to the cytokine receptor-like factor 3 family.,similarity:Contains 1 fibronectin type-III domain.,tissue specificity:Expressed in skin and squamous cell carcinoma (SCC) and in lesion actinic keratosis (AK),.

## Research Area

## Image Data



Western Blot analysis of HeLa lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

## Note

For research use only.