

## Summary

<b>Production Name</b>	CyPB Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	PPIB
<b>Alternative Names</b>	PPIB; CYPB; Peptidyl-prolyl cis-trans isomerase B; PPlase B; CYP-S1; Cyclophilin B; Rotamase B; S-cyclophilin; SCYLP
<b>Gene ID</b>	5479.0
<b>SwissProt ID</b>	P23284.The antiserum was produced against synthesized peptide derived from the C-terminal region of human PPIB. AA range:151-200

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000. ELISA: 1:20000.
<b>Molecular Weight</b>	24kD

## Background

**Product Name: CyPB Rabbit Polyclonal Antibody**  
**Catalog #: APRab09681**



The protein encoded by this gene is a cyclosporine-binding protein and is mainly located within the endoplasmic reticulum. It is associated with the secretory pathway and released in biological fluids. This protein can bind to cells derived from T- and B-lymphocytes, and may regulate cyclosporine A-mediated immunosuppression. Variants have been identified in this protein that give rise to recessive forms of osteogenesis imperfecta. [provided by RefSeq, Oct 2009], catalytic activity: Peptidylproline (omega=180) = peptidylproline (omega=0)., caution: It is uncertain whether Met-1 or Met-9 is the initiator., enzyme regulation: Cyclosporin A (CsA) inhibits CYPB., function: PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides., similarity: Belongs to the cyclophilin-type PPIase family. PPIase B subfamily., similarity: Contains 1 PPIase cyclophilin-type domain., subcellular location: Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,

## Research Area

## Image Data



Western Blot analysis of K562 cells using CyPB Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

## Note

For research use only.