

Summary

Production Name	GCNT4 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	GCNT4
Alternative Names	
Gene ID	51301.0
SwissProt ID	Q9P109.Synthesized peptide derived from human protein . at AA range: 60-140

Application

Dilution Ratio	WB 1:500-2000 ELISA 1:5000-20000
Molecular Weight	49kD

Background

catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,3-N-acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta-D-glucosaminyl-1,6)-N-acetyl-D-galactosaminyl-R.,function:Glycosyltransferase that mediates

Product Name: GCNT4 Rabbit Polyclonal Antibody Catalog #: APRab11361




core 2 O-glycan branching, an important step in mucin-type biosynthesis. Does not have core 4 O-glycan or I-branching enzyme activity.,online information:Core 2 beta-1,6-N-acetylglucosaminyltransferase 3,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 14 family,tissue specificity:Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.,catalytic activity:UDP-N-acetyl-Dglucosamine + beta-D-galactosyl-1,3-N-acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta-Dglucosaminyl-1,6)-N-acetyl-D-galactosaminyl-R,function:Glycosyltransferase that mediates core 2 O-glycan branching, an important step in mucin-type biosynthesis. Does not have core 4 O-glycan or I-branching enzyme activity.,online information:Core 2 beta-1,6-N-acetylglucosaminyltransferase 3,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 14 family,tissue specificity:Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.,

Research Area

O-Glycan biosynthesis;

Image Data



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

Note

For research use only.