

Summary

Production Name	G-CSFR Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	CSF3R
Alternative Names	CSF3R; GCSFR; Granulocyte colony-stimulating factor receptor; G-CSF receptor; G-CSF-R; CD114
Gene ID	1441.0
SwissProt ID	Q99062.The antiserum was produced against synthesized peptide derived from the Internal region of human CSF3R. AA range:321-370

Application

Dilution Ratio	WB 1:500-1:2000. ELISA: 1:20000.
Molecular Weight	92kD

Background

Product Name: G-CSFR Rabbit Polyclonal Antibody
Catalog #: APRab11371




The protein encoded by this gene is the receptor for colony stimulating factor 3, a cytokine that controls the production, differentiation, and function of granulocytes. The encoded protein, which is a member of the family of cytokine receptors, may also function in some cell surface adhesion or recognition processes. Alternatively spliced transcript variants have been described. Mutations in this gene are a cause of Kostmann syndrome, also known as severe congenital neutropenia. [provided by RefSeq, Aug 2010], alternative products: Additional isoforms seem to exist. Experimental confirmation may be lacking for some isoforms, disease: Defects in CSF3R may be a cause of severe congenital neutropenia (SCN) in some patients, domain: The box 1 motif is required for JAK interaction and/or activation, domain: The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding, function: Receptor for granulocyte colony-stimulating factor (CSF3). In addition it may function in some adhesion or recognition events at the cell surface, similarity: Belongs to the type I cytokine receptor family. Type 2 subfamily, similarity: Contains 1 Ig-like C2-type (immunoglobulin-like) domain, similarity: Contains 5 fibronectin type-III domains, subunit: Homodimer. The dimeric receptor binds two CSF3 molecules, tissue specificity: One or several isoforms have been found in myelogenous leukemia cell line KG-1, leukemia U937 cell line, in bone marrow cells, placenta, and peripheral blood granulocytes. Isoform GCSFR-2 is found only in leukemia U937 cells. Isoform GCSFR-3 is highly expressed in placenta,

Research Area

Cytokine-cytokine receptor interaction; Jak_STAT; Hematopoietic cell lineage; Pathways in cancer;

Image Data



Western Blot analysis of SKOV3 cells using G-CSFR Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

Note

For research use only.