

Summary

Production Name	Girdin Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	CCDC88A
	CCDC88A; APE; GRDN; KIAA1212; Girdin; Akt phosphorylation enhancer; APE; Coiled-
Alternative Names	coil domain-containing protein 88A; G alpha-interacting vesicle-associated protein;
	GIV; Girders of actin filament; Hook-related protein 1; HkRP1
Gene ID	55704.0
SwissProt ID	Q3V6T2.The antiserum was produced against synthesized peptide derived from human
	Girdin. AA range:1383-1432

Application

Dilution Ratio IHC 1:100-1:300 ELISA: 1:5000

Molecular Weight

Product Name: Girdin Rabbit Polyclonal Antibody Catalog #: APRab11447




Background

This gene encodes a member of the Girdin family of coiled-coil domain containing proteins. The encoded protein is an actin-binding protein that is activated by the serine/threonine kinase Akt and plays a role in cytoskeleton remodeling and cell migration. The encoded protein also enhances Akt signaling by mediating phosphoinositide 3-kinase (PI3K)-dependent activation of Akt by growth factor receptor tyrosine kinases and G protein-coupled receptors. Increased expression of this gene and phosphorylation of the encoded protein may play a role in cancer metastasis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011], function: Enhances phosphoinositide 3-kinase (PI3K)-dependent phosphorylation and kinase activity of AKT1/PKB, but does not possess kinase activity itself. Phosphorylation of AKT1/PKB thereby induces the phosphorylation of downstream effectors GSK3 and FOXO1/FKHR, and regulates DNA replication and cell proliferation (By similarity). Essential for the integrity of the actin cytoskeleton and for cell migration. Required for formation of actin stress fibers and lamellipodia. May be involved in membrane sorting in the early endosome.,PTM:Phosphorylation is induced by epidermal growth factor (EGF) in a phosphoinositide 3-kinase (PI3K)-dependent manner. Phosphorylation by AKT1/PKB is necessary for the delocalization from the cell membrane and for cell migration, sequence caution: Intron retention at the C-terminus, similarity: Belongs to the CCDC88 family.,subcellular location:Localizes to the cell membrane through interaction with phosphoinositides., subunit: Homodimer. The non-phosphorylated form interacts with phosphatidylinositol 4-phosphate [PI(4)P] and weakly with phosphatidylinositol 3-phosphate [PI(3)P]. Interacts with microtubules. Interacts with actin through its C-terminal domain. Interacts with the C-terminus of AKT1/PKB., tissue specificity: Expressed ubiquitously.,

Research Area

Image Data



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Girdin Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: Girdin Rabbit Polyclonal Antibody Catalog #: APRab11447



Note

For research use only.