

Summary

Production Name	$G\alpha$ t2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4° C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	GNAT2
Alternative Names	GNAT2; GNATC; Guanine nucleotide-binding protein G(t) subunit alpha-2; Transducin
	alpha-2 chain
Gene ID	2780.0
SwissProt ID	P19087.The antiserum was produced against synthesized peptide derived from human
	GNAT2. AA range:1-50

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:10000
Molecular Weight	40kD



Background

Transducin is a 3-subunit guanine nucleotide-binding protein (G protein) which stimulates the coupling of rhodopsin and cGMP-phoshodiesterase during visual impulses. The transducin alpha subunits in rods and cones are encoded by separate genes. This gene encodes the alpha subunit in cones. [provided by RefSeq, Jul 2008],disease:Defects in GNAT2 are the cause of achromatopsia type 4 (ACHM4) [MIM:139340]. Achromatopsia is an autosomal recessively inherited visual disorder that is present from birth and that features the absence of color discrimination.,function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. Transducin is an amplifier and one of the transducers of a visual impulse that performs the coupling between rhodopsin and cGMP-phosphodiesterase.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,tissue specificity:Retinal rod outer segment.,

Research Area





Image Data

Western blot analysis of lysates from COLO cells, using GNAT2 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from Jurkat cells using GNAT2 antibody.



Note For research use only.