

Summary

eceptor Rabbit Polyclonal Antibody
l Antibody
se

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	lgG
Clonality	Polyclonal
Form	Liquid
Stores	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
Storage	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	HRH1
Alternative Names	HRH1; Histamine H1 receptor; H1R; HH1R
Gene ID	3269.0
SwissProt ID	P35367.The antiserum was produced against synthesized peptide derived from human
	HRH1. AA range:141-190

Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.
Molecular Weight	60kD


Background

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. The protein encoded by this gene is an integral membrane protein and belongs to the G protein-coupled receptor superfamily. It mediates the contraction of smooth muscles, the increase in capillary permeability due to contraction of terminal venules, the release of catecholamine from adrenal medulla, and neurotransmission in the central nervous system. It has been associated with multiple processes, including memory and learning, circadian rhythm, and thermoregulation. It is also known to contribute to the pathophysiology of allergic diseases such as atopic dermatitis, asthma, anaphylaxis and allergic rhinitis. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by Reffunction:In peripheral tissues, the H1 subclass of histamine receptors mediates the contraction of smooth muscles, increase in capillary permeability due to contraction of smooth muscles, increase in capillary permeability and catecholamine release from adrenal medulla, as well as mediating neurotransmission in the central nervous system.,PTM:Potential sites of phosphorylation in the third cytoplasmic loop may play an important role in regulating signal transduction through the receptor molecule.,similarity:Belongs to the G-protein coupled receptor 1 family.,

Research Area


Calcium;Neuroactive ligand-receptor interaction;

Image Data




Immunofluorescence analysis of LOVO cells, using HRH1 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from COLO205 cells, using HRH1 Antibody. The lane on the right is blocked with the



Western Blot analysis of various cells using Histamine H1 Receptor Polyclonal Antibody diluted at 1: 2000

Note

For research use only.