**Antibody** 

Catalog #: APRab12139



# Summary

hnRNP C1/C2 Rabbit Polyclonal Antibody **Production Name** 

Description Rabbit Polyclonal Antibody

Rabbit Host

**Application** IF,IHC,WB,ELISA Reactivity Human, Mouse, Rat

# **Performance**

Conjugation Unconjugated Modification Unmodified

IgG Isotype

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

### **Immunogen**

**Gene Name HNRNPC** 

**Alternative Names** HNRNPC; HNRPC; Heterogeneous nuclear ribonucleoproteins C1/C2; hnRNP C1/C2

Gene ID 3183.0

P07910. The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

hnRNP C1/C2. AA range:241-290

# **Application**

**Dilution Ratio** 

WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in

other applications.

**Molecular Weight** 33kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

**Antibody** 

Catalog #: APRab12139



## **Background**

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene can act as a tetramer and is involved in the assembly of 40S hnRNP particles. Multiple transcript variants encoding at least two different isoforms have been described for this gene. [provided by RefSeq, Jul 2008], function: Binds pre-mRNA and nucleates the assembly of 40S hnRNP particles. Single HNRNPC tetramers bind 230-240 nucleotides. Trimers of HNRNPC tetramers bind 700 nucleotides. May play a role in the early steps of spliceosome assembly and pre-mRNA splicing. Interacts with poly-U tracts in the 3'-UTR or 5'-UTR of mRNA and modulates the stability and the level of translation of bound mRNA molecules., PTM: Phosphorylated on Ser-260 and Ser-299 in resting cells. Phosphorylated on Ser-253 and on 1 serine residue in the poly-Ser stretch at position 238 in response to hydrogen peroxide, PTM: Sumoylated. Sumoylation reduces affinity for mRNA., similarity: Belongs to the RRM HNRPC family. RALY subfamily., similarity: Contains 1 RRM (RNA recognition motif) domain., subcellular location: Component of ribonucleosomes., subunit: Tetramer composed of 3 copies of isoform C1 and 1 copy of isoform C2. Assembly of 3 tetramers with bound pre-mRNA gives rise to a 19S complex that interacts with HNRNPA2B1 tetramers. Component of the 40S hnRNP particle. Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRNPA1, HNRNPA2B1, HNRPA3, HNRNPC, HNRPF, HNRPH1, HNRPK, HNRPM, HNRNPR, HNRNPU, KIAA1160, KIAA1604, LSM2, LSM3, MAGOH, MORG1, PABPC1, PLRG1, PNN, PPIE, PPIL1, PPIL3, PPWD1, PRPF19, PRPF4B, PRPF6, PRPF8, RALY, RBM22, RBM8A, RBMX, SART1, SF3A1, SF3A2, SF3A3, SF3B1, SF3B2, SF3B3, SFRS1, SKIV2L2, SNRPA1, SNRPB, SNRPB2, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF, SNRPG, SNW1, SRRM1, SRRM2, SYF2, SYNCRIP, TFIP11, THOC4, U2AF1, WDR57, XAB2 and ZCCHC8.,

#### Research Area

Spliceosome;


### **Image Data**

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


**Antibody** 

Catalog #: APRab12139






Immunofluorescence analysis of HUVEC cells, using hnRNP C1/C2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using hnRNP C1/C2 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from HepG2 and COLO205 cells, using hnRNP C1/C2 Antibody. The lane on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab12139





Western blot analysis of the lysates from HT-29 cells using hnRNP C1/C2 antibody.



Western Blot analysis of various cells using hnRNP C1/C2 Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA) .

#### Note

For research use only.