
Summary

Production Name	IL-17C Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	IL17C UNQ561/PRO1122
Alternative Names	Interleukin-17C (IL-17C;Cytokine CX2)
Gene ID	27189.0
SwissProt ID	Q9P0M4.Synthetic peptide from human protein at AA range: 81-130

Application

Dilution Ratio	IHC-p 1:50-200, ELISA 1:10000-20000.
Molecular Weight	

Background

The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine


Product Name: IL-17C Rabbit Polyclonal Antibody
Catalog #: APRab12506




was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells. [provided by RefSeq, Jul 2008],function:Stimulates the release of tumor necrosis factor alpha and IL-1-beta from the monocytic cell line THP-1.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,

Research Area


Image Data



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:200



Product Name: IL-17C Rabbit Polyclonal Antibody
Catalog #: APRab12506



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:200

Note

For research use only.